arXiv:1506.07132v3 [math.SP] 24 Sep 2015

Spectral Statistics of Random Schrödinger Operators with Unbounded Potentials

Dhriti Ranjan Dolai*and Anish Mallick†

September 25, 2015

Abstract. In this work we investigate the spectral statistics of random Schrödinger operators $H^\omega = -\Delta + \sum_{n\in\mathbb{Z}^d}(1+|n|^\alpha)q_n(\omega)|\delta_n\rangle\langle\delta_n|$, $\alpha > 0$ acting on $\ell^2(\mathbb{Z}^d)$ where $\{q_n\}_{n\in\mathbb{Z}^d}$ are i.i.d random variables distributed uniformly on $[0,1]$.

AMS 2000 MSC: 35J10, 81Q10, 35P20.
Keywords: Anderson model, Poisson statistics, Unbounded potential.

1 Introduction

We consider the random Scrödinger operators with unbounded potentials given by

$$H^\omega = H_0 + \sum_{n\in\mathbb{Z}^d} (1+|n|^\alpha)\ q_n(\omega)\ |\delta_n\rangle\langle\delta_n| \qquad \text{with } \alpha > 1, \tag{1.1}$$

on the Hilbert space $\ell^2(\mathbb{Z}^d)$. The set $\{\delta_n : n \in \mathbb{Z}^d\}$ is the standard basis for the Hilbert space, $|n| = \left(\sum n_i^2\right)^{\frac{1}{2}}$, $\{q_n\}_n$ are i.i.d random variables distributed uniformly on $[0,1]$, and

$$\begin{aligned}(H_0\phi)(n) &= \sum_{|n-m|_+=1} \phi(n) - \phi(m), \quad \phi \in \ell^2(\mathbb{Z}^d), \quad \|H_0\| = 4d, \qquad (1.2)\\ &= 2d\ u(n) - \sum_{|n-m|_+=1} u(m) = 2du(n) - (\Delta u)(n).\end{aligned}$$

Where $|n|_+ = \sum |n_i|$. We realize $q_n(\omega)$ as a random field on $\Omega = \otimes_{n\in\mathbb{Z}^d}[0,1]$ and let $\mathbb{P}$ be the product probability measure on Ω. Henceforth we will use $V_n(\omega) = b_n q_n(\omega)$ where $b_n = (1+|n|^\alpha)$ and we write

$$V(\omega) = \sum_{n\in\mathbb{Z}^d} V_n(\omega)|\delta_n\rangle\langle\delta_n|,$$

so one can express H^ω as

$$H^\omega = 2d - \Delta + V(\omega). \tag{1.3}$$

The spectral theory for the operator (1.1) was studied by Gordon-Molchanov-Tsagani [18] for one-dimension and Gordon-Jakšić-Mochanov-Simon [17] for higher dimension.

We require some results from [17] to formulate our problem. Following their notation, set $C(X)$

---

*Institute: Chennai Mathematical Institute, India. email: drdolai@cmi.ac.in
†Institute: The Institute of Mathematical Sciences, India. email: anishm@imsc.res.in

for $X \subset \mathbb{Z}^d$ to be the set of all $\phi \in \ell^2(\mathbb{Z}^d)$ with supp $\phi \subseteq X$. The form $\mathscr{D}$ associated to H_0 is defined by

$$\mathscr{D}(\phi) = \sum_{\substack{(n,m) \\ |n-m|_+=1}} |\phi(n) - \phi(m)|^2, \quad \phi \in \ell^2(\mathbb{Z}^d). \tag{1.4}$$

Denote

$$\Lambda(X) = \inf_{\substack{\|\phi\|=1 \\ \phi \in C(X)}} \mathscr{D}(\phi), \tag{1.5}$$

$$a_k = \inf_{A_k} \Lambda(A_k), \tag{1.6}$$

where A_k are connected (there is a path between every two points) subsets of $\mathbb{Z}^d$ with $\#A_k = k$. Then $\{a_k\}_{k \geq 0}$ (set $a_0 = \infty$) is a strictly decreasing sequence of positive numbers. We denote $H^\omega_{\Lambda_L}$ to be the restriction of H^ω to the cube $\Lambda_L = \{(n_1, n_2, \cdots, n_d) : |n_i| \leq L\} \subset \mathbb{Z}^d$ and

$$N^\omega_L(E) := \#\big\{j : E_j \leq E,\ E_j \in \sigma(H^\omega_{\Lambda_L})\big\},\ E \in \mathbb{R}.$$

Following results are recalled from Gordon-Jakšić-Molchanov-Simon [17].

Theorem 1.1 ([17, Theorem 1.2]). *If $d/k \geq \alpha > d/(k+1)$ for positive integer k, for a.e ω*

(i) $\sigma(H^\omega) = \sigma_{pp}(H^\omega)$ and eigenfunctions of H^ω decay at least exponentially,

(ii) $\sigma_{ess}(H^\omega) = [a_k, \infty)$,

(iii) $\#\sigma_{disc}(H^\omega) < \infty$.

Theorem 1.2 ([17, Theorem 1.4]).

(i) If $d/k > \alpha > d/(k+1)$ and $E \in (a_j, a_{j-1}),\ 1 \leq j \leq k$, then

$$\lim_{L\to\infty} \frac{N^\omega_L(E)}{L^{d-j\alpha}} = N_j(E)$$

exist for a.e ω and is a non-random function.

(ii) If $\alpha = d/k$ and $E \in (a_j, a_{j-1})$, $1 \leq j < k$, the above is valid. If $E \in (a_k, a_{k-1})$ then

$$\lim_{L\to\infty} \frac{N^\omega_L(E)}{lnL} = N_k(E)$$

exists for a.e ω and is a non-random function.

In both cases, N_j is a continuous function on (a_j, a_{j-1}), and

$$N_j(E) \sim C_j(E - a_j)^j \quad as \quad E \to a_j.$$

The constants C_j are of combinatorial nature.

From (1.5) and (1.6) it is easy to see $a_1 \leq 2d$. In this work we investigated the eigenvalue statistics in the region (a_1, ∞). Associated with $H^\omega_{\Lambda_L}$ we define the point process

$$\xi^\omega_{L,E} = \sum_{x \in \sigma(H^\omega_{\Lambda_L})} \delta_{L^{d-\alpha}(x-E)}, \tag{1.7}$$

From theorem 1.2 it is clear that the average spacing between eigenvalues of $H^\omega_{\Lambda_L}$ in the region (a_1, ∞) is of order $L^{-(d-\alpha)}$, since $\lim\limits_{L\to\infty} \frac{N^\omega_L(E)}{L^{d-\alpha}}$ exists and non-random a.e ω. Theorem 1.1 make sure that the region under consideration has point spectrum.

Our goal is to show that the weak limit of the sequence of point processes $\{\xi^{\omega}_{L,E}\}_L$ is Poisson point process with intensity measure $N_1'(E)dx$. But for that we need some nice convergence and continuity of N_1'.

We will work with $\alpha > 1$, and will focus on $G \cap \mathbb{R}$ where the set G is given by

$$G = \{z \in \mathbb{C} : Rez > 2M^2, Imz > -d\} \setminus \cup_{n\in\mathbb{Z}^d}\{z : |z - (b_n + 2d)| < 2M^2\}. \tag{1.8}$$

where we choose $M > 2d$ (so that we can show analyticity of $\mathbb{E}^{\omega}[G(z; n, n)]$ in the set G as in lemma 3.3 below).

A portion of set G is shown in Figure 1. All the analysis is done in the connected component which contains the set $\{z : Imz, Rez > 2M^2\}$. It can be seen that the set $G \cap \mathbb{R}$ is union of intervals and the lengths are increasing as one moves left. On other hand this is not the case for $\alpha < 1$.



Figure 1

When $\alpha > 1$, the set $G \cap \mathbb{R}$ contain intervals of length greater than $4d$. So using remark 2.5 below, one can get the existence of E such that $N_1'(E) > 0$. Remark 3.6 provide the continuity of N_1' in $G \cap \mathbb{R}$ and so implies existence of intervals $\{I_n\}_n$ such that $\cup_n I_n \subset \{E : N_1'(E) > 0\}$. In each of these intervals I_n, Lemma 3.5 provides an uniformly convergence of $\{f_L\}$, the densities of ν_L (see (2.2)) to N_1'. Set $S = \cup_n I_n$ where the intervals $\{I_n\}_n$ as described above.
Now we are in the position to state our main result:

Theorem 1.3. *Let $E \in S$, then the sequence of point process $\{\xi^{\omega}_{L,E}\}_L$ converges weakly to the Poisson point process with intensity measure $N_1'(E)dx$ (Lebesgue measure). So for any bounded Borel set $B \in \mathbb{B}_{\mathbb{R}}$*

$$\lim_{L\to\infty} \mathbb{P}\big(\omega : \xi^{\omega}_{L,E}(B) = n\big) = e^{-N_1'(E)|B|}\frac{\big(N_1'(E)|B|\big)^n}{n!} \quad for \quad n = 0, 1, 2, \cdots, .$$

The study of eigenvalue statistics was done by Molchanov [31] in one-dimension and by Minami [32] for Anderson model at high disorder with stationary potentials (see also Germinet-Klopp [16]). In [14] Dolai-Krishna obtained Poisson statistics for discrete Anderson model with uniformly Hölder continuous single site potential. Eigenfunction statistics was studied by Nakano [34] for continuous model and Killip-Nakano [24] for lattice case with bounded density for single site distribution. Dolai-Mallick [30] studied the eigenfunction statistics for Hölder continuous single site distribution on lattice.

For decaying potentials (i.e $\alpha < 0$) the spectral statistics was studied on several occasions. Killip-Stoiciu [25] studied the CMV matrices whose matrix elements decay like $n^{-\alpha}$. They showed that, for (i) $\alpha > 1/2$ statistics is the clock, (ii) $\alpha = 1/2$ limiting process is circular $\beta-$ensemble, (iii) $0 < \alpha < 1/2$ the statistics is Poisson. Analogues of Killip-Stoiciu[25] was done by Kotani-Nakano[26] for the one-dimensional Schrödinger operator with decaying potentials in the continuum model and obtained the same statistics for $\alpha > 1/2$ and $\alpha = 1/2$. Krichevski-Valkó-Virág [28] studied the one-dimensional discrete Schrödinger operator with the random potential decaying like $n^{-1/2}$ and obtained the Sineβ-process. In [13] Dolai-Krishna consider the Anderson Model with decaying Random Potentials and shown that statistics inside $[-2d, 2d]$ for dimension $d \geq 3$ is independent of the randomness and agrees with that of the free part Δ. But to best of our knowledge this is the first occasion that spectral statistics was investigated for the model described by (1.1).

2 Preliminaries

For any $\Lambda \subset \mathbb{Z}^d$ we consider the canonical orthogonal projection χ_Λ onto $\ell^2(\Lambda)$ and define the matrices

$$H^\omega_\Lambda = \big(\langle \delta_n, H^\omega \delta_m\rangle\big)_{n,m\in\Lambda},\ G^\Lambda(z;n,m) = \langle \delta_n, (H^\omega_\Lambda - z)^{-1}\delta_m\rangle,\ G^\Lambda(z) = (H^\omega_\Lambda - z)^{-1}. \tag{2.1}$$

$$G(z) = (H^\omega - z)^{-1},\ \ G(z;n,m) = \langle \delta_n, (H^\omega - z)^{-1}\delta_m\rangle,\ \ z \in \mathbb{C}^+.$$

Note that H^ω_Λ is the matrix

$$\chi_\Lambda H^\omega \chi_\Lambda\ :\ \ell^2(\Lambda) \longrightarrow \ell^2(\Lambda),\ a.e\ \omega.$$

Define

$$\nu_L(\cdot) = \frac{1}{L^{d-\alpha}} \sum_{n\in\Lambda_L} \mathbb{E}^\omega\big(\langle \delta_n, E_{H^\omega_{\Lambda_L}}(\cdot)\delta_n\rangle\big), \tag{2.2}$$

where $\Lambda_L \subset \mathbb{Z}^d$ is the box centre at origin of side length $2L+1$. Note that

$$\sum_{n\in\Lambda_L} b_n^{-1} = \sum_{n\in\Lambda_L} (1+|n|^\alpha)^{-1} = \Theta(L^{d-\alpha}). \tag{2.3}$$

For any bounded interval $I=[a,b]$ we have

$$\nu_L(I) = \frac{1}{L^{d-\alpha}} \mathbb{E}^\omega[N^\omega_L(b) - N^\omega_L(a)].$$

In this section we calculate the Minami and Wegner estimate and show the positivity of the measure ν given by

$$N_1(x) = \nu(a_1, x),\ for\ x > a_1, \tag{2.4}$$

here N_1 is defined as in (i) of Theorem 1.2.
We start with spectral averaging estimate for the model (1.1).

Proposition 2.1. *Let $I=[a,b]\subset\mathbb{R}$ be a bounded interval and $\Lambda \subseteq \mathbb{Z}^d$. Consider $H^\omega_\Lambda = \chi_{\scriptscriptstyle\Lambda} H_0 \chi_{\scriptscriptstyle\Lambda} + V(\omega)\chi_{\scriptscriptstyle\Lambda}$ then*

$$\mathbb{E}^\omega(\langle \delta_n, E_{H^\omega_\Lambda}(I)\delta_n\rangle) \le \pi(1+|n|^\alpha)^{-1}|I|, \qquad n\in\Lambda.$$

Proof. Write H^ω_Λ as

$$\begin{aligned} H^\omega_\Lambda &= \left(\chi_{\scriptscriptstyle\Lambda} H_0 \chi_{\scriptscriptstyle\Lambda} + \sum_{n\neq k\in\Lambda} b_k q_k(\omega)|\delta_k\rangle\langle\delta_k|\right) + b_n q_n(\omega)|\delta_n\rangle\langle\delta_n| \\ &= H^{\omega/n}_\Lambda + b_n q_n(\omega)|\delta_n\rangle\langle\delta_n|. \end{aligned}$$

From resolvent equation one has

$$\langle \delta_n, (H^\omega_\Lambda - z)^{-1}\delta_n\rangle = \frac{1}{b_n q_n(\omega) + \langle \delta_n, (H^{\omega/n}_\Lambda - z)^{-1}\delta_n\rangle^{-1}}.$$

Take $z = \sigma + i\tau \in \mathbb{C}^+ = \{z\in\mathbb{C} : Imz > 0\}$ and set

$$\langle \delta_n, (H^{\omega/n}_\Lambda - z)^{-1}\delta_n\rangle^{-1} = c + id,\ A = b_n^{-1}\frac{c}{c^2+d^2}, B = b_n^{-1}\frac{d}{c^2+d^2} > 0,\ when\ Imz > 0.$$

The above estimate and the uniform distribution of q_n on $[0,1]$ gives

$$\int_{\mathbb{R}} Im\langle\delta_n,(H^\omega_\Lambda - z)^{-1}\delta_n\rangle d\mu(q_n) = \frac{1}{b_n}\int_{\mathbb{R}} \frac{B}{(x+A)^2+B^2} d\mu(x) \leq \pi b_n^{-1}.$$

The following is immediate from above,

$$\mathbb{E}^\omega\big(Im\langle\delta_n,(H^\omega_\Lambda - \sigma - i\tau)^{-1}\delta_n\rangle\big) \leq \pi b_n^{-1}. \tag{2.5}$$

Using Fubini theorem it follows

$$\mathbb{E}^\omega\Bigg(\int_a^b Im\langle\delta_n,(H^\omega_\Lambda - \sigma - i\tau)^{-1}\delta_n\rangle d\sigma\Bigg) \leq \pi b_n^{-1}|I|.$$

Using Stone's formula [36, Theorem VII.13]

$$\frac{\pi}{2}\mathbb{E}^\omega[\langle\delta_n E_{H^\omega_\Lambda}[a,b]\delta_n\rangle + \langle\delta_n E_{H^\omega_\Lambda}(a,b)\delta_n\rangle] \leq \pi b_n^{-1}|I|.$$

In particular $\mathbb{E}^\omega(\langle\delta_n E_{H^\omega_\Lambda}(\{r\})\delta_n\rangle) = 0$ for $r \in \mathbb{R}$. So

$$\mathbb{E}^\omega(\langle\delta_n, E_{H^\omega_\Lambda}(I)\delta_n\rangle) \leq \pi(1+|n|^\alpha)^{-1}|I|, \quad n \in \Lambda \subseteq \mathbb{Z}^d.$$

□

From the above proposition together with (2.2) and (2.3) we get:

Corollary 2.2. *For any interval I we get*

$$\nu_L(I) \leq C|I|, \tag{2.6}$$

where C is a constant independent of L.

So it is clear that ν_L is absolutely continuous measure, we denote it's density by f_L i.e $f_L = \frac{d\nu_L}{dx}$. Combes-Germinet-Klein [9] (see also Combes-Hislop-Klopp[8]) estimated the Wegner and Minami estimates for generalised Anderson Model. The following proposition give the Wegner and Minami estimate for the Model (1.1).

Proposition 2.3. *For $\Lambda \subset \mathbb{Z}^d$ finite volume and bounded interval $I \subset \mathbb{R}$ we have*

$$\mathbb{E}\big(TrE_{H^\omega_\Lambda}(I)\big) \leq \sum_{n\in\Lambda} b_n^{-1}|I| \tag{2.7}$$

$$\mathbb{E}\Bigg(\big(TrE_{H^\omega_\Lambda}(I)\big)\big(TrE_{H^\omega_\Lambda}(I)-1\big)\Bigg) \leq \Bigg(\sum_{n\in\Lambda} b_n^{-1}|I|\Bigg)^2. \tag{2.8}$$

The Wegner estimate (2.7) follows from Proposition 2.1. The proof of (2.8) goes exactly the same way as in [9, Theorem 2.1] once we have (2.7). So we omit the proof here.

Lemma 2.4. *If $a, b \in (4d,\infty) \subset (a_1,\infty)$ with $b - a > 4d$ then $N_1(b) - N_1(a) > 0$.*

Proof. Define

$$A^\omega_{L,0} = \sum_{n\in\Lambda_L} b_n q_n(\omega), \quad A^\omega_{L,4d} = 4d + \sum_{n\in\Lambda_L} b_n q_n(\omega),$$

$$N^\omega_{L,4d}(E) = \#\{j : E_j \leq E, \quad E_j \in \sigma(A^\omega_{L,4d})\}, \quad N^\omega_{L,0} = \#\{j : E_j \leq E, \quad E_j \in \sigma(A^\omega_{L,0})\}.$$

From (1.2) it is clear that $0 \leq H_0 \leq 4d$ which leads to following operator inequality

$$A^\omega_{L,0} \leq H^\omega_{\Lambda_L} \leq A^\omega_{L,4d}. \tag{2.9}$$

Using min-max principle we get

$$N^\omega_{L,4d}(E) \le N^\omega_L(E) \le N^\omega_{L,0}(E). \tag{2.10}$$

Let $E = 4d + \epsilon, \quad \epsilon > 0$ then we have

$$\begin{aligned} N^\omega_{L,4d}(E) &= \#\{n \in \Lambda_L : 4d + b_n q_n(\omega) \le 4d + \epsilon\} \\ &= \#\{n \in \Lambda_L : q_n(\omega) \le b_n^{-1}\epsilon\}. \end{aligned} \tag{2.11}$$

Taking expectation of previous equation

$$\begin{aligned} \mathbb{E}^\omega(N^\omega_{L,4d}(E)) &= \#\{n \in \Lambda_L : b_n^{-1}\epsilon \ge 1\} + \sum_{\substack{n\in\Lambda_L \\ b_n^{-1}\epsilon<1}} b_n^{-1}\epsilon \\ &= \sum_{n\in\Lambda_L} b_n^{-1}\epsilon + \#\{n \in \Lambda_L : b_n^{-1}\epsilon \ge 1\} - \sum_{\substack{n\in\Lambda_L \\ b_n^{-1}\epsilon\ge1}} b_n^{-1}\epsilon. \end{aligned} \tag{2.12}$$

Since last two terms of the above are bounded, so we get

$$\lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \mathbb{E}^\omega(N^\omega_{L,4d}(E)) = \epsilon \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \sum_{n\Lambda_L} b_n^{-1}. \tag{2.13}$$

Similar calculation for $A^\omega_{L,0}$ gives

$$\lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \mathbb{E}^\omega(N^\omega_{L,0}(E)) = (4d+\epsilon) \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \sum_{n\in\Lambda_L} b_n^{-1}. \tag{2.14}$$

Let $I = [a,b] \subset (4d,\infty) \subset (a_1,\infty)$ with $a = 4d + \epsilon_a,\ b = 4d + \epsilon_b$ and $\epsilon_b - \epsilon_a > 4d$, then we get

$$\begin{aligned} N_1(b) - N_1(a) &= \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} [N^\omega_L(b) - N^\omega_L(a)] \\ &= \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \mathbb{E}^\omega[N^\omega_L(b) - N^\omega_L(a)] \quad (\textit{using Theorem} 1.2\ (i)) \\ &\ge \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} [N^\omega_{L,4d}(b) - N^\omega_{L,0}(a)] \quad (\textit{using}(2.10)) \\ &= [\epsilon_b - \epsilon_a - 4d] \lim_{L\to\infty} \frac{1}{L^{d-\alpha}} \sum_{n\in\Lambda_L} (1+|n|^\alpha)^{-1} \quad (\textit{using}(2.13)\ \textit{and}\ (2.14)) \\ &= [\epsilon_b - \epsilon_a - 4d] C > 0.\ (\textit{using}\ (2.3)) \end{aligned} \tag{2.15}$$

□

Remark 2.5. *From second line of (2.15) and (2.6) we get, for any bounded interval* $I = [a,b] \subset (a_1,\infty)$

$$N_1(b) - N_1(a) \le C|I|.$$

In other words the measure ν *given by* $N_1(x) = \nu(a_1,x], \quad x \in (a_1,\infty)$ *is absolutely continuous. Now (i) of Theorem 1.2 gives*

$$\nu_L \xrightarrow[\textit{weakly}]{L\to\infty} \nu \quad \textit{on} \quad (a_1,\infty). \tag{2.16}$$

If $I \subset (4d,\infty)$ *is an interval with* $|I| > 4d$ *then the above lemma implies*

$$\nu(I) > 0, \tag{2.17}$$

and so there exists $E \in I$ *such that* $N_1'(E) > 0$.

3 Regularity of the density of ν_L

In this Section we used an idea from Kaminaga-Krishna-Nakamura [22] to prove regularity. In the large disorder case, the kinetic energy part of the Random Hamiltonian can be viewed as perturbation of diagonal randomness. In such case one can use a random walk expansion, in which the expectation of each of the terms turns out to have analytic continuation through real axis in a region. In case of growing randomness a similar idea works and the local density of states has an analytic continuation. The proves used here are modification of Kaminaga-Krishna-Nakamura [22].

For $\ell \in \mathbb{N}$, set

$$B_\ell(z) = \int_0^1 \frac{dx}{(ax-z)^\ell}, \quad a \geq 1, \quad z \in \mathbb{C}^+.$$

Proposition 3.1. *$B_\ell(z)$ has analytic continuation from $\mathbb{C}^+$ to $\mathbb{C} \setminus \{x+iy : x = 0, a \ \&y < 0\}$ through $\mathbb{R} \setminus \{0, a\}$, and inside the region of analyticity we have the estimates*

$$|B_\ell(z)| \leq \begin{cases} \frac{3\pi+1}{a} + \max\left\{\frac{1}{|z-a|}, \frac{1}{|Rez|}\right\} & \textit{if } \ell = 1, \\ \frac{1}{a}\left(\frac{1}{|a-z|^{\ell-1}} + \frac{1}{|Rez|^{\ell-1}}\right) & \textit{if } \ell > 1. \end{cases} \tag{3.1}$$



Proof. By explicit calculation

$$B_\ell(z) = \begin{cases} \frac{1}{a} \log \frac{z-a}{z} & \text{if } \ell = 1, \\ \frac{1}{a(\ell-1)}\left(\frac{1}{(a-z)^{\ell-1}} - \frac{1}{(-z)^{\ell-1}}\right) & \text{if } \ell > 1. \end{cases} \tag{3.2}$$

Above function is well defined and analytic in the region $\mathbb{C} \setminus \{x+iy : x = 0, a \ \& \ y < 0\}$.

On the domain $\mathbb{C} \setminus \{x+iy : x = 0, a \ \& \ y < 0\}$ we have $\frac{-\pi}{2} < arg(z), arg(z-a) \leq \frac{3\pi}{2}$. Using $|\log \frac{z_1}{z_2}| = |\log \frac{z_2}{z_1}|$, $\ln x < x$ for $x > 1$ and $\frac{z}{z-a} = 1 + \frac{a}{z-a}$, $\frac{z-a}{z} = 1 - \frac{a}{z}$ to estimate $B_1(z)$ (for $z \in G$) we have:

$$\begin{aligned} |B_1(z)| &\leq \frac{1}{a}\left(3\pi + \max\left\{\left|\frac{z}{z-a}\right|, \left|\frac{z-a}{z}\right|\right\}\right) \\ &\leq \frac{1}{a}\left(3\pi + 1 + a\max\left\{\left|\frac{1}{z-a}\right|, \left|\frac{1}{z}\right|\right\}\right) \\ &\leq \frac{3\pi+1}{a} + \max\left\{\frac{1}{|z-a|}, \frac{1}{|Rez|}\right\}. \end{aligned} \tag{3.3}$$

For $\ell > 1$ we have

$$\begin{aligned} |B_\ell(z)| &\leq \frac{1}{|a(\ell-1)|}\left(\frac{1}{|a-z|^{\ell-1}} + \frac{1}{|-z|^{\ell-1}}\right) \\ &\leq \frac{1}{a}\left(\frac{1}{|a-z|^{\ell-1}} + \frac{1}{|Rez|^{\ell-1}}\right). \end{aligned} \tag{3.4}$$

(3.3), (3.4) together with (3.2) give the proposition. □

A simple calculation provides

Corollary 3.2. *If $Re z > 2M^2$ and $|z-a| > 2M^2$ with $M > 1$ then*

$$|B_\ell(z)| \leq \begin{cases} \frac{3\pi+1}{a} + \frac{1}{2M^2} & \textit{if } \ell = 1, \\ \frac{1}{aM^k} & \textit{if } \ell > 1. \end{cases} \tag{3.5}$$

The above estimate also valid for the analytic continuation of $B_\ell(z)$ on G.

Lemma 3.3. *Let Λ be a connected subset of $\mathbb{Z}^d$, then $\mathbb{E}^\omega\big(G^\Lambda(z;n,n)\big)$ is analytic in the region G (defined in (1.8)).*

Proof. Denote the set of all paths lying in Λ of length k with both initial and end point at $n \in \Lambda$ by $\Gamma^k_\Lambda(n)(\subset \Lambda^{k+1} \subset (\mathbb{Z}^d)^{k+1}$; note that $\#\Gamma^k_\Lambda(n) \leq (2d)^{k+1})$. The Neumann series expansion of the resolvent for $|Imz| > 2d$ is

$$\begin{aligned} G^\Lambda(z;n,n) &= \langle \delta_n, (H^\omega_\Lambda - z)^{-1}\delta_n\rangle \qquad (3.6)\\ &= \langle \delta_n, \chi_\Lambda(2d - \Delta + V(\omega) - z)^{-1}\chi_\Lambda\delta_n\rangle \quad (\textit{see } (1.3))\\ &= \left\langle \delta_n, \left((V(\omega)+2d-z)^{-1}\sum_{k=0}^{\infty}(-1)^k\big(\chi_\Lambda\Delta\chi_\Lambda(V(\omega)+2d-z)^{-1}\big)^k\right)\delta_n\right\rangle \end{aligned}$$

Which can be expressed as Random walk expansion:

$$\begin{aligned} G^\Lambda(z;n,n) &= \sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{j=0}^{k}\big(V_{n_j}(\omega)+2d-z\big)^{-1} \qquad (3.7)\\ &= \sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{j=0}^{k}\big(b_{n_j}q_{n_j}(\omega)+2d-z\big)^{-1}. \end{aligned}$$

Here $\gamma = (n_0, n_1, \cdots, n_k) \in \Gamma^k_\Lambda(n)$ with $|n_j - n_{j-1}|_+ = 1, \quad j = 1, 2, \cdots, k$.
Denote $\#(\gamma, m) = \#\{j : n_j \in \gamma \textit{ and } n_j = m\}$ for $m \in \mathbb{Z}^d$; note that $\#(\gamma, m) = 0$ except for finitely many m, hence the products appearing below are finite. Also note that $\sum_{m\in\mathbb{Z}^d}\#(\gamma,m) = k+1$ for $\gamma \in \Gamma^k_\Lambda(n)$. Using independence of single site potentials in (3.7) gives

$$\begin{aligned} \mathbb{E}^\omega\big(G^\Lambda(z;n,n)\big) &= \mathbb{E}^\omega\Big[\sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{m\in\mathbb{Z}^d}\big(b_mq_m(\omega)+2d-z\big)^{-\#(\gamma,m)}\Big] \qquad (3.8)\\ &= \sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{m\in\mathbb{Z}^d}\int_0^1\frac{dx}{(b_mx+2d-z)^{\#(\gamma,m)}}\\ &= \int_0^1\frac{dx}{(b_nx+2d-z)} + \sum_{k=1}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{m\in\mathbb{Z}^d}\int_0^1\frac{dx}{(b_mx+2d-z)^{\#(\gamma,m)}} \end{aligned}$$

For $z \in G$ and using (3.5) and Proposition 3.1 we have

$$\begin{aligned} \left|\sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma^k_\Lambda(n)}\prod_{m\in\mathbb{Z}^d}\int_0^1\frac{dx}{(b_mx+2d-z)^{\#(\gamma,m)}}\right| &\leq \frac{3\pi+1}{b_n} + \frac{1}{M} + \sum_{k=1}^{\infty}\frac{1}{\prod\limits_m b_m}\frac{|\Gamma^k_\Lambda(n)|}{M^{\sum_{m\in\mathbb{Z}^d}\#(\gamma,m)}} \qquad (3.9)\\ &\leq \frac{3\pi+1}{b_n} + \frac{1}{M} + \frac{1}{b_n}\sum_{k=1}^{\infty}\left|\frac{2d}{M}\right|^{k+1}, \quad \big(\prod_m b_m \geq b_n\big)\\ &< \infty, \quad \textit{as} \quad M > 2d. \end{aligned}$$

The expression $\sum_{k=0}^{\infty}\sum_{\gamma\in\Gamma_{\Lambda}^{k}(n)}\prod_{m\in\mathbb{Z}^d}\int_0^1\frac{dx}{(b_m x+2d-z)^{\#(\gamma,m)}}$ is analytic in the region G (using above and [7, 6.2 Weierstrass M-test]) and it agrees with $\mathbb{E}^{\omega}\big(G^{\Lambda}(z;n,n)\big)$ for $Imz > 2d$. So it is analytic continuation of $\mathbb{E}^{\omega}\big(G^{\Lambda}(z;n,n)\big)$ in the region G.

□

Define

$$g_L(z) = \frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\mathbb{E}^{\omega}\Big[\big(G^{\Lambda_L}(z;n,n)\big) - \big\langle\delta_n, \big(V(\omega)+2d-z\big)^{-1}\delta_n\big\rangle\Big], \tag{3.10}$$

then g_L is analytic on G and the same estimate as we did for the second term of the RHS of (3.8) gives

$$\begin{aligned}|g_L(z)| &\leq \frac{C'}{L^{d-\alpha}}\sum_{n\in\Lambda_L} b_n^{-1}, \quad C' = \sum_{k=1}^{\infty}\left|\frac{2d}{M}\right|^{k+1}, \quad z\in G \\ &\leq C \qquad (using\ (2.3)).\end{aligned} \tag{3.11}$$

Above estimate gives the uniform bound of $g_L(z)$ on G.

Above proposition together with Montel's theorem (see [7, 2.9 Montel's Theorem]) gives uniform convergence on compact sets of G (also called compact convergence) hence the corollary.

Corollary 3.4. *Every subsequence of $\{g_L\}$ has a further subsequence which converges compactly on G.*

It is easy to see that $G\cap(4d,\infty)$ contains intervals I with $|I| > 4d$; so using (2.17) we can choose $E \in G\cap(4d,\infty)$ such that $N_1'(E) > 0$.

Lemma 3.5. *If we choose E as described above then $\{f_L\}$ (f_L is the density of ν_L) converges uniformly to N_1' on $\{x : |x-E| \leq \delta\} \subset \mathbb{R}$.*

Proof. Write

$$Img_L = Im\psi_L(z) - Im\phi_L(z), \tag{3.12}$$

where

$$\psi_L(z) = \frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\mathbb{E}^{\omega}\big(G^{\Lambda_L}(z;n,n)\big),$$

$$\begin{aligned}\phi_L(z) &= \frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\mathbb{E}^{\omega}\Big(\big\langle\delta_n, \big(V(\omega)+2d-z\big)^{-1}\delta_n\big\rangle\Big) \\ &= \frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\int_0^1\frac{1}{b_n x+2d-z}dx.\end{aligned}$$

Corollary 3.4 together with Lemma A.1 in Appendix give that every subsequence of $\{Im\psi_L\}$ has a further subsequence which converges compactly on $\{z\in\mathbb{C} : |z-E|\leq\delta\}\subset\mathbb{C}$.

But we can get the density of a measure through the imaginary part of it's Borel transform (see [37, (iv), Theorem 11.6]). So we have

$$f_L(x) = \frac{d\nu_L}{dx}(x)$$

$$= \frac{1}{\pi} Im\psi_L(x+i0).$$

Now we get that every subsequence of $\{f_L\}$ has a further subsequence which converges compactly on $\{x \in \mathbb{R} : |x-E| \leq \delta\}$. Now using (2.16)

$$f_L \xrightarrow[uniformly]{L\to\infty} N_1' \;\; on \;\; \{x : |x-E| \leq \delta\} \subset \mathbb{R}. \tag{3.13}$$

Complete the proof. □

Remark 3.6. *Analyticity of ψ_L implies continuity of $Im\psi_L$ in the region G. So the uniform convergence in* (3.13) *implies continuity of N_1'.*

4 Exponential decay of Green's function

Here we study the exponential decay for fractional moment of Green function using Aizenman-Molchanov [1] method for the model. To estimate exponential decay of $\mathbb{E}^\omega\big(|(G^B(z;n,m)|^s\big)$ we require the following lemma. The proof essentially follows the argument given in [1, Theorem II.1].

Lemma 4.1. *For any two pair of sites $n, m \in \mathbb{Z}^d$ we have*

$$\mathbb{E}^\omega\big(|(G^\Lambda(z;n,m)|^s\big) \leq C,$$

where C is a positive constant independent of n, m and B.

Proof. **Case-I ($n = m$):** From resolvent Identity we have

$$G^\Lambda(z;n,n) = \frac{1}{b_n q_n(\omega) + b} \;\;,$$

where b is the inverse of the resolvent for the operator $H^\omega_\Lambda - b_n q_n(\omega)|\delta_n\rangle\langle\delta_n|$ and is independent of q_n.

From above we get

$$\begin{aligned}\mathbb{E}^\omega(|G^\Lambda(z;n,n)|^s) &= \frac{1}{b_n^s}\mathbb{E}^\omega\bigg(\frac{1}{|q_n(\omega)+b_n^{-1}b)|^s}\bigg) \\ &= \frac{1}{b_n^s}\mathbb{E}^{\omega/n}\bigg(\int_0^1 \frac{d\mu(q_n)}{|q_n(\omega)+b_n^{-1}b)|^s}\bigg) \\ &\leq \frac{1}{b_n^s}\sup_{a\in\mathbb{C}}\int_0^1 \frac{dx}{|x-a|^s} \\ &\leq C < \infty.\end{aligned} \tag{4.1}$$

In the above we used the uniform distribution of q_n in $[0,1]$, $b_n \geq 1 \;\; \forall \;\; n \in \mathbb{Z}^d$ and $0 < s < 1$.

Case-II ($n \neq m$): From inequality II.7 of [1] with $\lambda = 1$ we have

$$\mathbb{P}\big(\omega : |(G^\Lambda(z;n,m)| \geq t\big) \leq \mathbb{P}\bigg(\omega : d_1\big(\big(V_n, V_m\big), S\big) \leq \frac{1}{t}\bigg). \tag{4.2}$$

Here we used notation (for d_1, V_n, V_m, S) as in [1, Theorem II.1].

Now using set equality's II.8 and inequality II.9 of [1] we get

$$\mathbb{P}\bigg(\omega : d_1\big(\big(V_n, V_m\big), S\big) \leq \frac{1}{t}\bigg) \leq \mathbb{P}\bigg(\big|V_n - S_1(n)\big| \leq \frac{\sqrt{2}}{t}\bigg) + \mathbb{P}\bigg(\big|V_m - S_1(m)\big| \leq \frac{\sqrt{2}}{t}\bigg) \tag{4.3}$$

$$
\begin{aligned}
&= \mathbb{P}\Big(\big|b_n q_n - S_1(n)\big| \le \frac{\sqrt{2}}{t}\Big) + \mathbb{P}\Big(\big|b_m q_m - S_2(m)\big| \le \frac{\sqrt{2}}{t}\Big)\\
&= \int_{b_n^{-1}(S_1-\frac{\sqrt{2}}{t})}^{b_n^{-1}(S_1+\frac{\sqrt{2}}{t})} d\mu(q_n) + \int_{b_n^{-1}(S_2-\frac{\sqrt{2}}{t})}^{b_n^{-1}(S_2+\frac{\sqrt{2}}{t})} d\mu(q_m)\\
&\le \Big(\frac{2\sqrt{2}}{b_n t} + \frac{2\sqrt{2}}{b_m t}\Big).
\end{aligned}
$$

Here we used the uniform distribution of μ on $[0,1]$ and $b_n = 1 + |n|^\alpha$.
Using (4.2) and (4.3)

$$
\mathbb{P}\big(\omega : |(G^\Lambda(z;n,m)| \ge t\big) \le \Big(\frac{2\sqrt{2}}{b_n t} + \frac{2\sqrt{2}}{b_m t}\Big). \tag{4.4}
$$

which gives us

$$
\begin{aligned}
\mathbb{E}^\omega\big(|(G^\Lambda(z;n,m)|^s\big) &= \int_0^\infty \mathbb{P}\big(\omega : |(G^\Lambda(z;n,m)|^s \ge t\big)\ dt \qquad (4.5)\\
&= \int_0^\infty \mathbb{P}\big(\omega : |(G^\Lambda(z;n,m)| \ge t^{1/s}\big)\ dt\\
&\le \int_0^\infty \min\Big(1, \frac{2\sqrt{2}}{b_n t^{1/s}}\Big)\ dt + \int_0^\infty \min\Big(1, \frac{2\sqrt{2}}{b_m t^{1/s}}\Big)\ dt\\
&\le C_1\big(b_n^{-1} + b_m^{-1}\big) \int_0^\infty \min\Big(1, \frac{2\sqrt{2}}{t^{1/s}}\Big)\ dt\\
&\le C'\big(b_n^{-1} + b_m^{-1}\big)\\
&= C'\Big[\frac{1}{1+|n|^\alpha} + \frac{1}{1+|m|^\alpha}\Big]\\
&\le 2C' = C < \infty.
\end{aligned}
$$

From (4.1) and (4.5) we conclude the proof. □

Lemma 4.2. *For any $\Lambda \subseteq \mathbb{Z}^d$, $z \in \mathbb{C}^+$, $0 < s < 1$ and $\beta > 0$ there exists $R_{\beta,s}, C_{\beta,s}$ such that for $|n-m| > R_{\beta,s}$*

$$
\mathbb{E}^\omega\big(|(G^\Lambda(z;n,m)|^s\big) \le C_{\beta,s} e^{-\beta|n-m|}, \quad r > 0. \tag{4.6}
$$

Proof. Using resolvent equation we have

$$
G_\omega^\Lambda(z;n,m) = \frac{\sum_{\substack{|n-p|=1\\ p\in\Lambda}} G^\omega_{\Lambda\setminus\{n\}}(z;p,m)}{b_n q_n(\omega) - z - a_n^\omega},
$$

where

$$
a_n^\omega = \sum_{\substack{p,q\in\Lambda\\ |n-p|=1\\ |n-q|=1}} G^\omega_{\Lambda\setminus\{n\}}(z;p,q).
$$

Observe that a_n^ω are independent of $q_n(\omega)$. This is the first step for the self-avoiding walk (SAW) representation of Green's function (see [12, Section 4] for more details). This give

$$
\mathbb{E}^\omega[|G_\omega^\Lambda(z;n,m)|^s] = \mathbb{E}^\omega\left[\left|\frac{\sum_{\substack{|n-p|=1\\ p\in\Lambda}} G^\omega_{\Lambda\setminus\{n\}}(z;p,m)}{b_n q_n(\omega) - z - a_n^\omega}\right|^s\right]
$$

$$\leq \mathbb{E}^{\omega|q_n(\omega)}\left[\int_0^1 \frac{dq_n(\omega)}{|b_nq_n(\omega)-z-a_n^\omega|^s}\sum_{\substack{|n-p|=1\\ p\in\Lambda}}|G^\omega_{\Lambda\backslash\{n\}}(z;p,m)|^s\right].$$

Using the inequality

$$\int_0^1\frac{dx}{|x-z|^s}<\frac{3-s}{1-s}=C_s \qquad \forall \;\; z\in\mathbb{C},$$

we have

$$\begin{aligned}
\mathbb{E}^\omega[|G^\Lambda_\omega(z;n,m)|^s] &\leq \frac{C_s}{b_n^s}\sum_{\substack{p\in\Lambda\backslash\{n\}\\ |n-p|=1}}\mathbb{E}^\omega[|G^\omega_{\Lambda\backslash\{n\}}(z;p,m)|^s]\\
&\leq \sum_{\gamma\in\Gamma(n,k)}\left(\prod_{i=0}^{k}\frac{C_s}{b^s_{\gamma_i}}\right)\left(\sum_{\substack{p\in\Lambda\backslash\{\gamma_0,\cdots,\gamma_k\}\\ |\gamma_k-p|=1}}\mathbb{E}^\omega[|G^\omega_{\Lambda\backslash\{\gamma_0,\cdots,\gamma_k\}}(z;p,m)|^s]\right)\\
&\leq 2dC'\sum_{\gamma\in\Gamma(n,|n-m|)}\prod_{i=0}^{|n-m|}\frac{C_s}{b^s_{\gamma_i}}\leq 2dC'\sum_{\gamma\in\Gamma(n,|n-m|)}e^{|n-m|\ln C_s-s\alpha\sum_{i=0}^{|n-m|}\ln|\gamma_i|},
\end{aligned}$$

where C' is the constant coming from lemma 4.1 and $\Gamma(n,k)$ are self avoiding paths (i.e for any $\gamma\in\Gamma(n,k)$ and $i\neq j$ we have $\gamma_i\neq\gamma_j$) starting from n of length k. Choose $R=\left(2dC_se^\beta\right)^{\frac{1}{\alpha s}}$, and choose n,m such that $|n-m|>R^{2d}$. Then using the inequality $\ln b_n\geq\max\{\alpha\ln|n|,0\}$ we have

$$\begin{aligned}
|n-m|\ln(2dC_s)-s\alpha\sum_{i=0}^{|n-m|}\ln|\gamma_i| &\leq |n-m|\ln(2dC_s)-s\alpha\sum_{\substack{i=0\\ |\gamma_i|>R}}^{|n-m|}\ln|\gamma_i|\\
&\leq |n-m|\ln(2dC_s)-(|n-m|-R^d)\ln\left(2dC_se^\beta\right)\\
&= -\beta|n-m|+R^d\ln\left(2dC_se^\beta\right).
\end{aligned}$$

Combining both of them:

$$\mathbb{E}^\omega[|G^\Lambda_\omega(z;n,m)|^s]\leq 2dC'(2dC_se^\beta)^{R^d}e^{-\beta|n-m|}.$$

So setting $R_{\beta,s}=\left(2dC_se^\beta\right)^{\frac{2d}{\alpha s}}$, we get a constant $C_{s,\beta}$ such that for $|n-m|>E_{\beta,s}$ we have

$$\mathbb{E}^\omega[|G^\Lambda_\omega(z;n,m)|^s]\leq C_{s,\beta}e^{-\beta|n-m|}.$$

□

Remark 4.3. *From the Lemma 4.2 together with Simon-Wolff criterion [37, Theorem 12.5] it follows that for almost all* ω *the entire spectrum of* H^ω *is pure point, i.e* $\sigma(H^\omega)=\sigma_{pp}(H^\omega)$ *a.e* ω.

5 Proof of Theorem 1.3

Here we exploit Minami's ([32]) idea to prove the main result. Divide Λ_L into N_L^d numbers of disjoint cubes C_p, $(p = 1, 2, \cdots, N_L^d)$ with side length $\frac{2L+1}{N_L}$, i.e

$$\Lambda_L = \sqcup_p C_p \quad with \ \ |C_p| = \left(\frac{2L+1}{N_L}\right)^d. \tag{5.1}$$

Define

$$\partial C_p = \{n \in C_p : \exists\ n' \in \mathbb{Z}^d \setminus C_p \ \ such\ that\ |n - n'| = 1\}$$
$$int(C_p) = \{n \in C_p : dist(n, \partial C_p) > l_L\}.$$

here we take $l_L = \delta\ lnL$, $\ \delta > 0$ large enough and $N_L \approx (2L+1)^\epsilon$, $\ 0 < \epsilon < 1$. Define the point process $\eta^\omega_{L,p}$ associated with $H^\omega_{C_p}$ as

$$\eta^\omega_{L,p,E} = \sum_{x \in \sigma\left(H^\omega_{C_p}\right)} \delta_{L^{d-\alpha}(x-E)}. \tag{5.2}$$

Note that the matrices $\{H^\omega_{C_p}\}_p$ are statistically independent. the proof follows in two steps:

Step 1: In this step we show that $\{\xi^\omega_L\}$ and $\{\sum_p \eta^\omega_{L,p,E}\}$ converges to same limit in the sense of distribution. For this one has to verify:

$$\lim_{L\to\infty} \mathbb{E}^\omega\left\{\left|e^{-\int f d\xi^\omega_{L,E}} - e^{-\sum_p \int f d\eta^\omega_{L,p,E}}\right|\right\} = 0 \ \ for \ \ f \in C_c(\mathbb{R}). \tag{5.3}$$

The linear combination of the functions $\phi_z(x) = Im\frac{1}{x-z}$, $\ z \in \mathbb{C}^+$ are dense in $L^1(\mathbb{R})$ (for more details we refer to [19, Appendix: The Stone-Weierstrass Gavotte]). Also note that $|e^{-x} - e^{-y}| \leq |x - y|$, $\ x, y \geq 0$. So to prove (5.3), it is sufficient to prove

$$\lim_{L\to\infty} \mathbb{E}^\omega\left\{\left|\int \phi_z d\xi^\omega_{L,E} - \sum_p \int \phi_z d\eta^\omega_{L,p,E}\right|\right\} = 0. \tag{5.4}$$

Since Λ_L is the disjoint union of C_p $(p = 1, 2, \cdots, N_L^d)$, we have (denote $z_L = E + L^{-(d-\alpha)}z$)

$$\begin{aligned}\int \phi_z d\xi^\omega_{L,E} - \sum_p \int \phi_z d\eta^\omega_{L,p,E} &= \frac{1}{L^{d-\alpha}}\left[Tr\Big(ImG^{\Lambda_L}(z_L)\Big) - \sum_p Tr\Big(ImG^{C_p}(z_L)\Big)\right] \\ &= \frac{1}{L^{d-\alpha}}\sum_p \sum_{n\in C_p}\Big[ImG^{\Lambda_L}(z_L; n, n) - ImG^{C_p}(z_L; n, n)\Big].\end{aligned} \tag{5.5}$$

For $z \in \mathbb{C}^+$ and $n \in int(C_p)$, the resolvent equation gives us

$$G^{\Lambda_L}(z; n, n) - G^{C_p}(z; n, n) = \sum_{(m,k)\in\partial C_p} G^{\Lambda_L}(z; k, n)\ G^{C_p}(z; n, m),$$

here $(m, k) \in \partial C_p$ means $m \in \partial C_p$, $k \in \mathbb{Z}^d \setminus C_p$ with $|m - k| = 1$. Using the above perturbation formula in (5.5) we have

$$\begin{aligned}\left|\int \phi_z d\xi^\omega_{L,E} - \sum_p \int \phi_z d\eta^\omega_{L,p,E}\right| &\leq \frac{1}{L^{d-\alpha}}\sum_p \sum_{n\in C_p\setminus int(C_p)}\left[ImG^{\Lambda_L}(z_L; n, n) + ImG^{C_p}(z_L; n, n)\right] \\ &\quad + \frac{1}{L^{d-\alpha}}\sum_p \sum_{n\in int(C_p)} \sum_{(m,k)\in\partial C_p}\left|G^{\Lambda_L}(z_L; k, n)\right|\left|G^{C_p}(z_L; n, m)\right|\end{aligned} \tag{5.6}$$

$$= A_L + B_L.$$

estimating A_L and B_L as follows

$$\begin{aligned}
\mathbb{E}^{\omega}(A_L) &\leq \frac{2}{L^{d-\alpha}} \sum_{p} \sum_{n \in C_p \setminus int(C_p)} b_n^{-1}, \quad (see (2.5)) \qquad (5.7)\\
&= O\Big(L^{-(d-\alpha)} \Big(\frac{2L+1}{N_L}\Big)^{d-1-\alpha} N_L^{d-\alpha} l_L\Big), \quad \big(b_n = 1 + |n|^{\alpha}\big)\\
&= O\big(N_L L^{-1} l_L\big).\\
&= O\big(L^{-(1-\epsilon)} lnL\big), \quad N_L = (2L+1)^{\epsilon}, \quad 0 < \epsilon < 1.
\end{aligned}$$

and

$$\mathbb{E}^{\omega}(B_L) \leq \frac{1}{L^{d-\alpha}} \sum_{p} \sum_{n \in int(C_p)} \sum_{(m,k) \in \partial C_p} \big|G^{\Lambda_L}(z_L; k, n)\big|^{s} \, \big|G^{\Lambda_L}(z_L; k, n)\big|^{(1-s)} \, \big|G^{C_p}(z_L; n, m)\big|, \quad 0 < s < 1 \qquad (5.8)$$

Since $n \in int(C_p)$ and $(m, k) \in \partial C_p$ so we have $|n - k| > l_L > 2R$ for large enough L. So Proposition 4.2 gives

$$\mathbb{E}^{\omega}\big(\big|G_L^{\Lambda}(z_L; k, n)\big|^{s}\big) \leq Ce^{-rl_L}.$$

Using above estimate in (5.8) we have

$$\mathbb{E}^{\omega}(B_L) \leq \frac{C}{L^{d-\alpha}|Imz_L|^{2-s}} N_L^d \Big(\frac{2L+1}{N_L}\Big)^{d} \Big(\frac{2L+1}{N_L}\Big)^{d-1} l_L e^{-rl_L}.$$

Using $N_L = (2L+1)^{\epsilon}$, $0 < \epsilon < 1$, $l_L = \delta lnL$ $\delta > 0$ and $Imz_L = L^{-(d-\alpha)}z$ in above we get

$$\mathbb{E}^{\omega}(B_L) = O\big(L^{-\tilde{\delta}} lnL\big), \qquad (5.9)$$

where $\tilde{\delta} = r\delta - [2d - 1 + (1-s)(d-\alpha) + \epsilon(1-d)] > 0$ with

$$\delta > \frac{1}{r}[2d - 1 + (1-s)(d-\alpha) + \epsilon(1-d)].$$

Now we conclude (5.4) from (5.7), (5.9) and (5.6).

Step 2: In this step we show that $\sum_{p=1}^{N_L^d} \eta_{L,p,E}$ converge weakly to Poisson point process with intensity $N_1'(E)dx$. First we show that the triangular array $\{\eta_{L,p,E} : p = 1, 2, \cdots, N_L^d\}_L$ is uniformly asymptotically negligible. For this one need to show

$$\lim_{L \to \infty} \sup_{p} \mathbb{P}\big(\eta_{L,p,E}^{\omega}(I) \geq 1\big) = 0 \qquad (5.10)$$

for any bounded interval I (see [15, equation (11.2.2)]).

To show $\sum_{p=1}^{N_L^d} \eta_{L,p,E}$ converges weakly to Poisson point process we us [15, Theorem 11.2.V]. For that we need to verify:

$$\sum_{p=1}^{N_L^d} \mathbb{P}\big(\eta_{L,p,E}^{\omega}(I) \geq 2\big) = 0, \qquad as \;\; L \to \infty \qquad (5.11)$$

and

$$\sum_{p=1}^{N_L^d} \mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq 1\big) = N_1'(E)|I|, \quad as \quad L \to \infty. \tag{5.12}$$

For any bounded interval I.

The condition (5.10) is obtained using Wegner estimate (2.7) as follows:

$$\begin{aligned}\mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq 1\big) &\leq \mathbb{E}^{\omega}\big[\eta^{\omega}_{L,p,E}(I)\big] \\ &= \mathbb{E}^{\omega}\big[TrE_{H^{\omega}_{C_p}}\big(E + L^{-(d-\alpha)}I\big)\big] \\ &= \frac{|I|}{L^{d-\alpha}} \sum_{n \in C_p} (1 + |n|^{\alpha})^{-1} \\ &= O\bigg(L^{-(d-\alpha)}\Big(\frac{2L+1}{N_L}\Big)^{d-\alpha}\bigg) \\ &= O(N_L^{-(d-\alpha)}).\end{aligned}$$

So (5.10) is immediate.

The following expression is very useful to verify (5.11) and (5.12).

$$\mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq 1\big) = \mathbb{E}^{\omega}\big[\eta^{\omega}_{L,p,E}(I)\big] - \sum_{j \geq 2} \mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq j\big). \tag{5.13}$$

The Minami estimate (2.8) give the estimate for the second term of RHS in the above equation.

$$\begin{aligned}\sum_{j \geq 2} \mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq j\big) &= \sum_{j \geq 2} (j-1)\mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) = j\big) \qquad (5.14) \\ &\leq \sum_{j \geq 2} j(j-1)\mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) = j\big) \\ &= \mathbb{E}^{\omega}\big[\eta^{\omega}_{L,p,E}(I)\big(\eta^{\omega}_{L,p,E}(I) - 1\big)\big] \\ &= \mathbb{E}^{\omega}\Big[TrE_{H^{\omega}_{C_p}}\big(E + L^{-(d-\alpha)}I\big)\Big\{TrE_{H^{\omega}_{C_p}}\big(E + L^{-(d-\alpha)}I\big) - 1\Big\}\Big] \\ &= \bigg(\sum_{n \in C_p} (1 + |n|^{\alpha})^{-1} \frac{|I|}{L^{d-\alpha}}\bigg)^2 \quad (see\ (2.8)) \\ &= O\bigg(\bigg(\Big(\frac{2L+1}{N_L}\Big)^{d-\alpha} L^{-(d-\alpha)}\bigg)^2\bigg) \\ &= O\Big(N_L^{-2(d-\alpha)}\Big).\end{aligned}$$

The above estimate immediately give

$$\sum_{p} \sum_{j \geq 2} \mathbb{P}\big(\eta^{\omega}_{L,p,E}(I) \geq j\big) = O\Big(N_L^{-(d-\alpha)}\Big). \tag{5.15}$$

Since LHS of (5.11) is dominated by the LHS of above, so we conclude (5.11).

Now (5.4) also holds for characteristic function, so using Lemma 3.5 we get

$$\begin{aligned}\lim_{L\to\infty}\sum_{p}\mathbb{E}^{\omega}\big[\eta^{\omega}_{L,p,E}(I)\big] &= \lim_{L\to\infty}\mathbb{E}^{\omega}\big[\xi^{\omega}_{L,E}(I)\big] \qquad (5.16)\\ &= \lim_{L\to\infty}\mathbb{E}^{\omega}\Big(TrE_{H^{\omega}_{\Lambda_L}}\big(E+L^{-(d-\alpha)}I\big)\Big)\\ &= \lim_{L\to\infty}L^{d-\alpha}\nu_L\big(E+L^{-(d-\alpha)}\big) \quad (\textit{see } (2.2))\\ &= \lim_{L\to\infty}L^{d-\alpha}\int_{E+L^{-(d-\alpha)}I}f_L(x)dx\\ &= \lim_{L\to\infty}\int_I f_L\big(E+L^{-(d-\alpha)}y\big)dy\\ &= N_1'(E)|I|, \quad (\textit{see } (3.13)).\end{aligned}$$

Using (5.15) and (5.16) in (5.13) we get (5.12) completing the proof. ☐

A Appendix

Define

$$\phi_L(z)=\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\int_0^1\frac{dx}{b_nx-z_d},\quad z_d=z-2d$$

and

$$C=\lim_{L\to\infty}\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{1}{1+|n|^{\alpha}}\quad (\textit{see } 2.3).$$

Note that $G\subset\{z: Rez_d>0\}\setminus\{(1+|n|^{\alpha}):n\in\mathbb{Z}^d\}$.

Lemma A.1. *$Im\phi_L(z)\xrightarrow{L\to\infty}-CIm\ln(-z_d)$ uniformly over compact sets in the domain $\{z: Rez_d>0\}\setminus\{b_n+2d:n\in\mathbb{Z}^d\}$.*

Proof. By explicit computation one have:

$$\begin{aligned}\phi_L(z)&=\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\int_0^1\frac{dx}{(1+|n|^{\alpha})x-z_d}\\ &=\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{-\ln(-z_d)}{1+|n|^{\alpha}}+\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{\ln(1+|n|^{\alpha})}{1+|n|^{\alpha}}+\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{1}{1+|n|^{\alpha}}\ln\left(1-\frac{z_d}{1+|n|\alpha}\right)\end{aligned}$$

We need $Im\phi_L$ which is

$$Im\phi_L(z_d)=-Im\ln(-z_d)\left(\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{1}{|n|^{\alpha}}\right)+\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{1}{1+|n|^{\alpha}}Im\ln\left(1-\frac{z_d}{1+|n|^{\alpha}}\right) \tag{A.1}$$

estimating the second term of RHS in the above .

$$\left|\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L}\frac{1}{1+|n|^{\alpha}}\ln\left(1-\frac{z_d}{1+|n|^{\alpha}}\right)\right|\le\frac{1}{L^{d-\alpha}}\left|\sum_{n\in\Lambda_{|z_d|^{\frac{1}{\alpha}}}}\frac{1}{1+|n|^{\alpha}}\ln\left(1-\frac{z_d}{1+|n|^{\alpha}}\right)\right| \tag{A.2}$$

$$+\frac{1}{L^{d-\alpha}}\left|\sum_{n\in\Lambda_L\setminus\Lambda_{|z_d|^{\frac{1}{\alpha}}}}\frac{1}{1+|n|^\alpha}\ln\left(1-\frac{z_d}{1+|n|^\alpha}\right)\right|.$$

Note that on a compact subset of the given domain

$$C(z)=\left|\sum_{n\in\Lambda_{|z_d|^{\frac{1}{\alpha}}}}\frac{1}{1+|n|^\alpha}\ln\left(1-\frac{z_d}{1+|n|^\alpha}\right)\right|\leq M \tag{A.3}$$

where M is independent of L, and so $\frac{C(z)}{L^{d-\alpha}}$ goes to zero uniformly on compact subsets of the given domain.

We observed that on a compact set $\frac{|z_d|}{1+|n|^\alpha}<1$ all most all $n\in\mathbb{Z}^d$ but finitely many n. Next using $|\ln(1-z)|<|z|(1-|z|)^{-1}$ for $|z|<1$ we get the following estimate on a compact subset of the given domain.

$$\begin{aligned}\frac{1}{L^{d-\alpha}}\left|\sum_{n\in\Lambda_L\setminus\Lambda_{|z|^{\frac{1}{\alpha}}}}\frac{1}{1+|n|^\alpha}\ln\left(1-\frac{z_d}{1+|n|^\alpha}\right)\right| &\leq\frac{1}{L^{d-\alpha}}\sum_{n\in\Lambda_L\setminus\Lambda_{|z_d|^{\frac{1}{\alpha}}}}\frac{|z_d|}{(1+|n|^\alpha)^2}\left(1-\frac{|z_d|}{1+|n|^\alpha}\right)^{-1} \qquad \text{(A.4)}\\ &=O\big(L^{-(d-\alpha)}L^{d-2\alpha}\big)\\ &=O(L^{-\alpha}).\end{aligned}$$

Now using (A.2), (A.3) and (A.4) in (A.1) we get

$$Im\phi_L(z)\xrightarrow[uniformly]{L\to\infty}-CIm\ln(-z_d).$$

The above convergence is uniform on compact subsets of

$$\{z:Rez_d>0\}\setminus\{(1+|n|^\alpha):n\in\mathbb{Z}^d\}.$$

□

Acknowledgement: The authors are indebted to Prof Krishna Maddaly for go through the proofs carefully and making valuable comments.

References

[1] Aizenman, Michael; Molchanov, Stanislav: *Localization at large disorder and at extreme energies: an elementary derivation*, Commun. Math. Phys. **157(2)**, 245-278, 1993.

[2] Aizenman, Michael; Warzel, Simone: *The Canopy Graph and Level Statistics for Random Operators on Trees*, Mathematical Physics, Analysis and Geometry, **9(4)**, 291-333, 2006.

[3] Aizenman, Michael; Schenker, Jeffrey H; Friedrich, Roland M; Hundertmark, Dirk: *Finite-volume fractional-moment criteria for Anderson localization*, Commu. Math. Phys. **224 (1)**, 219-253, 2001.

[4] Böcker, Stefan; Kirsch, Werner; Stollmann, *Peter Spectral theory for nonstationary random potentials. Interacting stochastic systems*, 103-117, Springer, Berlin, 2005.

[5] Bourgain, Jean; Klein, Abel: *Bounds on the density of states for Schrödinger operators*, Invent. Math. **194(1)** (2013), 41-72.

[6] Carmona, René; Lacroix, Jean: *Spectral theory of random Schrödinger operators*, Boston, Birkhauser, 1990.

[7] Conway, J B: *Functions of One Complex variable*, Springer-Verlag, New York, 1973.

[8] Combes, Jean-Michel; Hislop, Peter D; Klopp, Frédéric: *An optimal Wegner estimate and its application to the global continuity of the integrated density of states for random Schrödinger operators*, Duke Math. J. **140(3)** (2007), 469-498.

[9] Combes, Jean-Michel; Germinet, François; Klein, Abel: *Generalized Eigenvalue-Counting Estimates for the Anderson Model*, J Stat Physics **135(2)**, 201-216, 2009.

[10] Combes, Jean-Michel; Germinet, François; Klein, Abel: *Poisson statistics for eigenvalues of continuum random Schrödinger operators*, J Anal. PDE **3(1)**, 49-80 (2010).

[11] Demuth, Michael; Krishna, M: *Determining Spectra in Quantum Theory*, Progress in Mathematical Physics. **44**, Birkhäuser, Boston, 2004.

[12] Dirk Hundertmark. *A short introduction to Anderson localization.* In *Analysis and stochastics of growth processes and interface models*, pages 194–218. Oxford Univ. Press, Oxford, 2008.

[13] Dolai, Dhriti; Krishna, M: *Level Repulsion for a class of decaying random potentials*, Markov Processes and Related Fields (to appear), arXiv:1305.5619[math.SP].

[14] Dolai, Dhriti; Krishna, M: *Poisson Statistics for Anderson Model with Singular Randomness*, Journal of the Ramanujan Mathematical Society (to appear), arXiv: 1408.4251[math.SP].

[15] Daley, D.J; Vere-Jones : *An Introduction to the Theory of Point Processes* **II**, *General theory and structure*, Springer, New York, 2008.

[16] Germinet, François; Klopp, Frédéric: *Spectral statistics for the discrete Anderson model in the localized regime*, Spectra of random operators and related topics, 11-24, RIMS Kkyroku Bessatsu, B27, Res. Inst. Math. Sci. (RIMS), Kyoto, 2011.

[17] Gordon, Y. A; Jakšić, V; Molchanov, S; Simon, B: *Spectral properties of random Schrödinger operators with unbounded potentials*, Comm. Math. Phys. **157(1)**, 23-50, 1993.

[18] Gordon, A. Y; Mochanov, S.A; Tsagani, B: *Spectral theory of one-dimensional Schrödinger operators with strongly fluctuation potential*, Funct. Anal Appl. **236(25)**, 1992.

[19] H. L. Cycon; R. G. Froese; W. Kirsch; B. Simon: *Schrödinger operators, with application to quantum mechanics and global geometry*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1987.

[20] Jakšić, Vojkan; Last, Yoram: *Spectral structure of Anderson type Hamiltonians*, Invent. Math, **141(3)**, 561-577, 2000.

[21] Kato, Tosio: *Perturbation theory for Linear operators*, Classics in Mathematics. Springer-Verlag, Berlin, 1995.

[22] Kaminaga, M; Krishna, M; Nakamura, S. *A note on the analyticity of density of states*, J. Stat. Phys. **149(3)**, 496-504, 2012.

[23] Kirsch, Werner: *An Invitation to Random Schrödinger Operators, (With an appendix by Frédéric Klopp)* Panor. Synthèses, **25**, Random Schrödinger operators, **1** 1-119, Soc. Math. France, Paris, 2008.

[24] Killip, Rowan; Nakano, Fumihiko: *Eigenfunction Statistics in the Localized Anderson Model*, Ann. Henri Poincare **8(1)**, 27-36, 2007.

[25] Killip, R; Stoiciu, M,: *Eigenvalue statistics for CMV matrices : from Poisson to clock via random matrix ensembles*, Duke Math. **146(3)**, 2009.

[26] Kotani, S; Nakano, Fumihiko: *Level statistics of one-dimensional Schrödinger operators with random decaying potential*,Spectral and scattering theory and related topics, 1924, RIMS Kôkyûroku Bessatsu, B45, Res. Inst. Math. Sci. (RIMS), Kyoto, 2014.

[27] Kirsch, W; Krishna, M; Obermeit, J: *Anderson model with decaying randomness: mobility edge*, Math.Z. **235(3)**, 421-433, 2000.

[28] Kritchevski, E; Valkó, B; Virág, B : *The scaling limit of the critical one- dimensional random Sdhrödinger operators*, Commun. Math. Phys. **334**, 775-806, 2012.

[29] Kotani, S; Ushiroya, N: *One-dimensional Schrödinger operators with random decaying potentials*, Commun. Math. Phys. **115(2)**, 247-266, 1988.

[30] Mallick, Anish; Dolai, Dhriti: *Eigenfunction Statistics for Anderson Model with Hölder continuous single site Potential*, arXiv:1409.7424 [math.SP].

[31] Molchanov, S.A: *The local structure of the spectrum of the one-dimensional Schrodinger operator*, Commun. Math. Phys. **78**, 429-446, 1981.

[32] Minami, Nariyuki: *Local Fluctuation of the Spectrum of a Multidimensional Anderson Tight Binding Model*, Commun. Math. Phys. **177(3)**, 709-725, 1996.

[33] Nakano, Fumihiko: *The repulsion between localizatgion centers in the Anderson model*, J. Stat. Phys. **123**, 803-810, 2006.

[34] Nakano, Fumihiko: *Innite divisibility of random measures associated to some random Schrödinger operators*, Osaka J. Math, **46**, 845-862, 2009.

[35] Pastur, Leonid; Figotin, Alexander: *Spectra of Random and Almost-Periodic Operators* Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 297. Springer-Verlag, Berlin, 1992. viii+587 pp.

[36] Reed, Michael; Simon, Barry: *Method of modern mathematical physics* **I**, *Functional Analysis*, Academic Press, 1978.

[37] Simon, Barry: *Trace ideals and their applications*, Mathematical Surveys and Monographs, 120, American Mathematical Society, Providence, RI, 2005. viii+150 pp.

[38] Stolz, Günter: *An introduction to the mathematics of Anderson localization. Entropy and the quantum II*, 71-108, Contemp. Math. 552, Amer. Math. Soc., Providence, RI, 2011.

[39] Simon, Barry; Wolff, Tom: *Singular continuos spectrum under rank one perturbations and localization for random Hamiltonians*, Comm. Pure and Appl. Math. **39(1)**, 75-90, 1986.

[40] Veselić, I: *Lipschitz-continuity of the integrated density of states for Gaussian random potentials* Lett. Math. Phys. **9 7**, 25-27, 2011.